

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>JUL 2011</b>		2. REPORT TYPE		3. DATES COVERED	
4. TITLE AND SUBTITLE <b>Analyzing The &amp;#948; Sco Binary In Anticipation Of A Disk-star Collision</b>		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>U.S. Naval Observatory, 3450 Massachusetts Avenue, N.W., Washington, DC, 20392</b>		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited.</b>					
13. SUPPLEMENTARY NOTES <b>The original document contains color images.</b>					
14. ABSTRACT <b>A current investigation is underway into the possible collision between a circumstellar disk and the secondary star in the &amp;#948; Scorpii binary system. &amp;#948; Scorpii is a prime candidate for a disk-star collision since the primary star has a circumstellar disk and the secondary star has a highly elliptical orbit with a period of approximately 10.5 years making the periastron passage very close to the primary star. The Navy Prototype Optical Interferometer (NPOI) was used to spatially resolve the two stars as well as the circumstellar disk around the primary. A revised orbit with new orbital parameters has been calculated using observations obtained between 2005 and 2010. Our results indicate periastron passage will occur on 2011-07-03.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES <b>2</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

# Analyzing the $\delta$ Sco binary in anticipation of a disk-star collision

Ashley Ames<sup>1</sup>, Christopher Tycner<sup>1</sup> and Robert Zavala<sup>2</sup>

<sup>1</sup>Dept. of Physics, Central Michigan University

<sup>2</sup>USNO, Flagstaff Station

**Abstract.** A current investigation is underway into the possible collision between a circumstellar disk and the secondary star in the  $\delta$  Scorpii binary system.  $\delta$  Scorpii is a prime candidate for a disk-star collision since the primary star has a circumstellar disk and the secondary star has a highly elliptical orbit with a period of approximately 10.5 years making the periastron passage very close to the primary star. The Navy Prototype Optical Interferometer (NPOI) was used to spatially resolve the two stars as well as the circumstellar disk around the primary. A revised orbit with new orbital parameters has been calculated using observations obtained between 2005 and 2010. Our results indicate periastron passage will occur on 2011-07-03.

**Keywords.** instrumentation: high angular resolution, astrometry, binaries: general

## 1. Introduction

$\delta$  Scorpii (HD143275, HR5953, FK5 0594) is a well known binary system with a highly eccentric orbit and a period of 10.5 years. The primary star is classified as a Be star with a gaseous circumstellar disk and the secondary is a B2-type star (Tango *et al.* 2009). The goal for this project is to refine the orbit of the secondary with respect to the primary and test for the possibility of a disk-star collision. The  $\delta$  Sco binary system was observed with the NPOI for a total of 108 nights. Data on two nights were obtained in July 2000 and the rest were obtained from June 2005 to August 2010.

## 2. Experimental Data

All the raw observational data, based on 108 nights, have been reduced using a standard NPOI reduction pipeline. The astrometric information is extracted from the reduced data giving the position of the secondary with respect to the primary. The NPOI observations were compared to orbits based on parameters from Mason *et al.* (2009), Tango *et al.* (2009), and Miroshnichenko *et al.* (2001). The new astrometric data, combined with radial velocities from Miroshnichenko *et al.* (2001), were used to calculate a refined orbit (shown in Figure 1) with the revised binary parameters listed in Table 1. Since the instrument used has a wide range of baselines, the resulting narrow-angle astrometry has high precision. In combination with extensive orbital coverage, including observations close to the previous periastron passage results in the best orbital parameters to date.

## 3. Summary

The binary orbit has been refined using astrometric data obtained from NPOI and radial velocities from Miroshnichenko *et al.* (2001). The orbit that is obtained gives a better fit to the data than previous findings. The next periastron passage date has been


**Table 1.** The orbital elements for  $\delta$  Sco.

Element	Mason <i>et al.</i> (2009)	Miroshnichenko <i>et al.</i> (2001)	Tango <i>et al.</i> (2009)	This Work
Period(y)	$10.68 \pm 0.05$	$10.58^1$	$10.74 \pm 0.02$	$10.81 \pm 0.002$
Semimajor axis (mas)	$104 \pm 6$	$107^1$	$98.3 \pm 1.2$	$99.0 \pm 0.05$
inclination, $i$ (deg)	$39 \pm 8$	$38 \pm 5$	$38 \pm 6$	$30.3 \pm 0.24$
Long. of asc. node, $\Omega$ (deg)	$153 \pm 9$	$175^1$	$175.2 \pm 0.6$	$172.3 \pm 0.6$
eccentricity, $e$	$0.94^2$	$0.94 \pm 0.01$	$0.9401 \pm 0.0002$	$0.941 \pm 0.0007$
Long. periastron, $\omega$	$29 \pm 12$	$-1 \pm 5$	$1.9 \pm 0.1$	$2.6 \pm 0.7$
T (Epoch of Periastron)	$J2000.693^2$	$J2000.693 \pm$ 0.008	$J2000.69389 \pm$ 0.00007	$J2000.6942 \pm$ 0.0011

Notes:

<sup>1</sup> Parameter adopted from Hartkopf *et al.* (1996) solution.

<sup>2</sup> Parameter adopted from Miroshnichenko *et al.* (2001) solution.



**Figure 1.** Binary orbit based on orbital parameters obtained from NPOI data. The black circles are the data points and the locations of the primary is marked with an X.

calculated to be 2011-07-02,  $22h \pm 1.1d$ . Our results indicate the periastron passage will occur 24 days later than the prediction of Tango *et al.* (2009).

## Acknowledgements

The Navy Prototype Optical Interferometer is a joint project of the Naval Research Laboratory and the US Naval Observatory, in cooperation with Lowell Observatory, and is funded by the Office of Naval Research and the Oceanographer of the Navy. A.A. and C.T. would also like to thank the Physics Department at Central Michigan University for support.

## References

- Mason, B. D., Hartkopf, W. I., Gies, D. R., Henry, T. J. *et al.* 2009, *AJ*, 137, 3358  
Hartkopf, W. I., Mason, B. D., & McAlister, H. A. 1996, *AJ*, 111, 370  
Miroshnichenko, A. S., Fabregat, J., Bjorkman, K. S., Knauth, D. C. *et al.* 2001, *A&A*, 377, 485  
Tango, W. J., Davis, J., Jacob, A. P., Mendez, A. *et al.* 2009, *MNRAS*, 396, 842